



AENSI Journals

Australian Journal of Basic and Applied Sciences

ISSN:1991-8178

Journal home page: www.ajbasweb.com



Screening for Enzymatic Activities in the Degradation Pathway of Homocholine by Soil Microorganisms

^{1,2,3}Isam A. Mohamed Ahmed, ^{1,2}Mohamed M. Eltayeb, ²Jiro Arima, ²Nobuhiro Mori, ³Norikazu Yamanaka and ³Takeshi Taniguchi

¹Department of Food Science and Technology, Faculty of Agriculture, University of Khartoum, Khartoum, Shambat 13314, Sudan.

²Department of Agricultural, Biological, and Environmental Sciences, Faculty of Agriculture, Tottori University, Tottori 680-8553, Japan.

³Arid Land Research Center, Tottori University, Tottori 680-0001, Japan.

ARTICLE INFO

Article history:

Received 23 December 2013

Received in revised form 25

February 2014

Accepted 26 February 2014

Available online 15 March 2014

Keywords:

Homocholine Microbial degradation

β -alanine betaine

Trimethylaminopropionaldehyde

ABSTRACT

Background: Although extensive research has been conducted on the microbial degradation of choline and its structurally related compounds, no research has been done on homocholine, a compound that resemble choline in many aspects of cholinergic metabolisms. The study of homocholine degradation enzymes may has potential for the improvement of the stress resistance of plant by introducing an efficient biosynthetic pathway for β -alanine betaine in genetically engineered economically relevant crop plant. **Objective:** This research was conducted to investigate the enzymatic activities in the degradation pathway of homocholine by bacterial strains isolated from the natural soil. **Results:** Screening for homocholine oxidation activities showed that NAD^+ -dependent dehydrogenase enzymes are predominant in all strains. Furthermore, dried cell reaction of *Pseudomonas* sp. strain A9 cells with homocholine in the presence and absence of NAD^+ demonstrated that the enzymes responsible for the metabolism of homocholine are dehydrogenases that require NAD^+ as electron acceptor. Moreover, in the cell free extract of *Pseudomonas* sp. strain A9 an inducible homocholine dehydrogenase activity was detected. In addition, an inducible NAD^+ -dependent 3-hydroxypropionate dehydrogenase activity was also detected in the cell free extract of *Pseudomonas* sp. strain A9. **Conclusion:** In *Pseudomonas* sp. strain A9, homocholine is oxidized to trimethylaminopropionaldehyde by homocholine dehydrogenase and, consequently, trimethylaminopropionaldehyde oxidized to β -alanine betaine by a NAD^+ -dependent aldehyde dehydrogenase. Thereafter, cleavage of β -alanine betaine C-N bond yielded trimethylamine and 3-hydroxypropionate. Thereafter, 3-hydroxypropionate was further oxidized to malonate semi-aldehyde by a NAD^+ -dependent 3-hydroxypropionate dehydrogenase.

© 2014 AENSI Publisher All rights reserved.

To Cite This Article: Isam A. Mohamed Ahmed, Mohamed M. Eltayeb, Jiro Arima, Nobuhiro Mori, Norikazu Yamanaka and Takeshi Taniguchi., Screening for Enzymatic Activities in the Degradation Pathway of Homocholine by Soil Microorganisms. *Aust. J. Basic & Appl. Sci.*, 8(2): 222-233, 2014

INTRODUCTION

Enzymes are remarkable catalysts capable of accepting a wide array of complex molecules as substrates and catalyzing reactions with unparalleled chiral (enantio-) and positional (regio-) selectivities (Schmid *et al.*, 2001). Such high selectivity also affords efficient reactions with few by-products, thereby making enzymes an environmentally friendly alternative to conventional chemical catalysts (Schmid *et al.*, 2001). Among various classes of enzymes, oxidoreductases represent a highly versatile class of biocatalysts for specific reduction and oxidation reactions, and currently used for the production of a wide variety of chemical and pharmaceutical products (Johannes *et al.*, 2006). To this decade, more than 1,000 oxidoreductases are known from which 80% use NADH as cofactor, 10% use the corresponding phosphates. The glucose-methanol-choline (GMC) oxidoreductase superfamily is composed of a group of enzymes that are able to catalyze the conversion of alcohols to the corresponding aldehydes or ketones. This family includes alcohol dehydrogenase, glucose oxidase, cholesterol oxidase, cellobiose dehydrogenase, pyranose 2-oxidase, choline dehydrogenase, choline oxidase and betaine aldehyde dehydrogenase (Cavener, 1992). The later three enzymes play an important role in the metabolism of choline in both mammalian and microorganisms. In choline degradation pathway, choline is oxidized to glycine betaine by two reactions: (i) choline to betaine aldehyde by either choline oxidase or choline dehydrogenase (Nagasawa *et al.*, 1976; 1997; Russell *et al.*, 1998) and (ii) betaine aldehyde to glycine betaine by a NAD^+ -dependent betaine aldehyde dehydrogenase (Mori *et al.*, 1992; Mori *et al.*, 1980; Mori *et al.*, 2001;

Corresponding Author: Isam A. Mohamed Ahmed, Department of Food Science and Technology, Faculty of Agriculture, University of Khartoum, Khartoum North P.O. Box 32, Shambat 13314, Sudan.
E-mail: isamnawa@yahoo.com , Tel: +249966896438 Fax: +24985318919.

Nagasawa *et al.*, 1976). Moreover, TMA-Butanol, a choline analogue, had a similar pathway as choline. In oxidative reaction, TMA-Butanol is converted to TMABaldehyde by a NAD⁺-dependent TMA-butanol dehydrogenase. Consequently, TMABaldehyde is oxidized to γ -butyrobetaine by a NAD⁺-dependent TMABaldehyde dehydrogenase (Hassan, 2008).

Homocholine (3-*N*-trimethylamino-1-propanol), an analogue of choline in which the amino alcohol group is lengthened by one CH₂-group, resembles choline in many aspects of cholinergic metabolism (Boksa and Collier, 1980). After transportation into rat brain synaptosome, it is acetylated and released as acetylhomocholine from a superior cervical ganglion and the mouse forebrain by a calcium-dependent process during depolarization (Collier *et al.*, 1977; Carroll and Aspary, 1980). It is also effective in preventing fat infiltration in both fat and cholesterol affected fatty livers (Channon *et al.*, 1937). Despite its similarity to choline in many aspects, reports on microorganisms degrading homocholine are scarce (Mohamed Ahmed *et al.* 2009a, b; Mohamed Ahmed *et al.* 2010) and consequently, its catabolic pathway remains unclear. Based on its structural similarity to choline, one would expect that homocholine could be degraded in a similar way and at a rate comparable to that of choline. It was found that, an osmoprotectant β -alanine betaine was an intermediate product in the degradation pathway of homocholine (Mohamed Ahmed *et al.*, 2009a, b; Mohamed Ahmed *et al.*, 2010). Therefore, the study of homocholine degradation pathways and the enzymes involved in them is of considerable interest. Using metabolomics, we previously postulated the degradation pathway of homocholine by strains belongs to the genera *Arthrobacter*, *Rhodococcus* and *Pseudomonas* (Mohamed Ahmed *et al.*, 2009a & b; Mohamed Ahmed *et al.*, 2010). Similar to choline and TMA-Butanol, homocholine was consequently oxidized to TMAPaldehyde and β -alanine betaine. Thereafter, cleavages of β -alanine betaine C-N bond yielded trimethylamine and alkyl chain. However, enzymatic activities in the degradation pathway of homocholine by isolated strains were not yet carried out. Therefore, the aim of the present study was to investigate the enzymatic activities in the degradation pathway of homocholine by the isolated strains and particularly *Pseudomonas* sp. strain A9.

MATERIALS AND METHODS

Materials:

3-Dimethylamino-1-propanol (DMA-Propanol), 4-dimethylamino-1-butanol (DMA-Butanol), and 4-aminobutanol were purchased from Tokyo Kassai (Tokyo, Japan). Homocholine iodide was prepared from 3-dimethylamino-1-propanol according to the method described previously (Mohamed Ahmed *et al.*, 2009a). TMAPaldehyde iodide was prepared from 3-aminopropionaldehyde dimethylacetal according to the method described previously (Mohamed Ahmed *et al.*, 2009a). Protease inhibitor cocktail Set II was from Calbiochem (Darmstadt, Germany; Table 4.1). Standard proteins for gel filtration were from BIO-RAD (Hercules, USA). All other reagents were commercial products of analytical grade from Wako Pure Chemicals Co. Ltd. (Tokyo, Japan).

Screening for the enzymatic activities by plate replica staining method:

In this method, the isolated strains (30 highly growth strains) were screened for homocholine oxidation activities using plate replica staining methods. The isolated strains were transferred individually using sterilized tooth stick to a single basal-homocholine agar plate medium containing, per liter, 10 g homocholine (HC), 2 g KH₂PO₄, 2 g K₂HPO₄, 0.5 g MgSO₄ · 7H₂O, 0.5 g yeast extract, 1 g polypeptone, and 15 g agar (pH 7.0). The plate was incubated at 30°C for either 24 or 48 h. The grown colonies were replicated onto a filter paper, treated with 1 ml of lysing solution (0.5% lysozyme and 10 mM EDTA in 100 mM potassium phosphate buffer, pH 8.0) and incubated at 37°C for 30 min. After lysis of the cells, the filter paper was sunk in a solution containing inhibitors of energy-generating system (10 mM of NaN₃, 10 mM of NaF, 1 mM of Na₂AsO₄) for 5 min. Then, the filter paper was frozen and thawed three times to inactivate the energy-generating system. After drying, the filter paper was sunk in 2 ml of activity staining solution containing 100 mM of glycine-NaOH buffer (pH 9.5), 64 μ M of 1-methoxy phenazine methosulfate, 0.24 mM of nitroblue tetrazolium (NBT), 1 mM of NAD⁺ and either 2 mM homocholine iodide or 1 mM of TMAPaldehyde iodide. The reduction of the NBT lead to the formation of the purple color is an indicator of the enzymatic activity. Strains showed positive result with replica staining were selected and used for preparation of cell free extract to confirm the enzyme activity by the spectrophotometric methods.

Cell free extract preparation:

To check the enzyme activities of the isolated strains, the strains were inoculated into 75 ml of basal-homocholine liquid medium and incubated at 25°C for 2 days in reciprocal shaker. At the end of cultivation period, bacterial cells were harvested by centrifugation at 10,000 \times g for 20 min and washed with 0.85 % KCl. The harvested bacterial cells were suspended in 30 ml of 0.2 M potassium phosphate buffer, pH 7.5, containing 1mM dithiothreitol (DTT) and disrupted by either sonication at 4~11°C for 15 min with 1.5 min run intervals or

by glass bead (0.1 mm) on dry ice for 10 cycles with 30 sec for each cycle. Cell free extract was obtained by centrifugation at $10,000 \times g$ for 20 min as supernatant and assayed for dehydrogenase and oxidase activities.

Homocholine dehydrogenase activities assays:

NAD⁺-dependent homocholine dehydrogenase activity was measured as an increase in the absorbance at 340 nm at 30°C. The standard reaction mixture (1.5 ml) contained 225 μmol of Tris-HCl buffer (pH 7.5), 33.3 μmol of homocholine iodide, 3 μmol of NAD⁺ and an appropriate amount of the enzyme. The reaction was started by the addition of NAD⁺ and the enzyme activity was calculated using an extinction coefficient of $6,200 \text{ M}^{-1} \cdot \text{cm}^{-1}$ of NADH at 340 nm. One unit of enzyme activity was defined as the amount of enzyme that catalyzes the formation of 1 μmol of NADH per min under the assay conditions. Specific activity was defined as units of enzyme activity per mg protein.

Membrane-bound homocholine dehydrogenase activity was measured as a decrease in the absorbance at 600 nm at 30°C using phenazine methosulfate (PMS)-2,6-dichlorophenol-indophenol (DCIP) assay system, according to the method of Nagasawa *et al.* (1976). The assay mixture (1.5 ml) contained 67.5 μmol of potassium phosphate buffer (pH 8.0), 1.5 μmol of potassium cyanide, 0.375 μmol of 1-methoxy PMS, 0.15 μmol of DCIP, 22.5 μmol of homocholine iodide and an appropriate amount of enzyme. The assay was started by the addition of homocholine iodide. The activity was calculated using an extinction coefficient of $21,500 \text{ M}^{-1} \cdot \text{cm}^{-1}$ for oxidized DCIP at 600 nm. One unit of enzyme activity is defined as the amount of enzyme that catalyzes the reduction of 1 μmole of DCIP per min under the standard assay conditions.

TMAPaldehyde dehydrogenase activity assay:

The standard reaction mixture (1.5 ml) contained 225 μmol of potassium phosphate buffer (pH 8.0), 15 μmol of TMAPaldehyde iodide, 3 μmol of NAD⁺, and an appropriate amount of the enzyme. The reaction was started by the addition of NAD⁺ and the enzyme activity was calculated using an extinction coefficient of $6,200 \text{ M}^{-1} \cdot \text{cm}^{-1}$ of NADH at 340 nm. One unit of enzyme activity was defined as the amount of enzyme that catalyzes the formation of 1 μmol of NADH per min under the assay conditions. Specific activity was defined as units of enzyme activity per mg protein.

3-Hydroxypropionate dehydrogenase activity assay:

3-Hydroxypropionate dehydrogenase activity was measured as an increase in the absorbance at 340 nm and at 30°C. The standard reaction mixture (1.5 ml) contained 225 μmol of Tris-HCl buffer (pH 7.5), 33.3 μmol of 3-hydroxypropionate, 3 μmol of NAD⁺ and an appropriate amount of the enzyme. The reaction was started by the addition of NAD⁺ and the enzyme activity was calculated using an extinction coefficient of $6,200 \text{ M}^{-1} \cdot \text{cm}^{-1}$ of NADH at 340 nm. One unit of enzyme activity was defined as the amount of enzyme that catalyzes the formation of 1 μmol of NADH per min under the assay conditions. Specific activity was defined as units of enzyme activity per mg protein.

Time course of the growth and enzyme formation of Pseudomonas sp. strains A9:

In order to determine the cultivation time needed for high enzyme activity formation, time course experiment was performed. Seed culture was prepared by cultivation of *Pseudomonas sp.* strain A9 in 5 ml basal-homocholine medium at 25°C for 24h with reciprocal shaking at 122 strokes/min. Then the turbidity of the seed culture was determined by measuring the optical density at 660 nm (Novaspec II, Amersham Pharmacia Biotech, Sweden). At the growth of 2.0 ($T_{660} = 2.0$), 200 μl of the seed culture broth was inoculated in 300 ml of basal-homocholine medium. Cultivation was carried out at 25°C on a reciprocal shaker. At intervals of 12, 15, 18, 22, and 26 h the growth was estimated and the bacterial cells were collected by centrifugation at $6,000 \times g$ (HITACHI HIMAC SCR 18B rotor: RPR9-2) for 30 min at 4°C and washed three times with 0.85% KCl. The harvested bacterial cells were suspended in 10 ml of 0.1 M potassium phosphate buffer, pH 7.5, containing 1mM dithiothreitol (DTT) and 1 mM EDTA, and disrupted by glass bead (0.1 mm) on dry ice for 5 min. Cell free extract was obtained by centrifugation at $10,000 \times g$ for 30 min as a supernatant and assayed for homocholine dehydrogenase activity.

Stability of homocholine dehydrogenase activity of Pseudomonas sp. strain A9:

In this method, cell free extract of each selected strain was dialyzed against potassium phosphate buffer, Tris-HCl buffer and Triethanolamine buffer (50 mM, pH 7.5) containing 1 mM of DTT and 1mM EDTA for overnight at 4°C. The dialyzed extracts were stored at 4°C for 0 to 10 days and the remaining enzyme activity was estimated. The stability of the enzyme against storage was also evaluated. The cell free extracts were stored at 4°C for 0 to 10 days and the remaining activity was measured. The stability of homocholine dehydrogenase was also examined in the presence of SH protecting reagents such as DTT, 2-mercaptoethanol and glutathione, as well as in the presence of stabilizing reagents such as ethanol, glycerol and ethylene glycol.

Formation of homocholine dehydrogenase on various media:

Pseudomonas sp. strain A9 was grown on basal medium supplemented with 1% of quaternary ammonium compounds such as homocholine, choline and TMA-butanol as a sole source of carbon and nitrogen. The strain was also cultivated on basal-homocholine medium supplemented with either glucose (1%) or glycerol (1%) as a carbon source or ammonium sulfate (0.5%) as a nitrogen source. Cultivation was carried out in 300 ml medium/1L cultural flask on a reciprocal shaker (120 strokes/min) at 25°C for 24h.

Intact and dry cell reaction of *Pseudomonas* sp. strain A9:

To further confirm the metabolic route of homocholine in *Pseudomonas* sp. strain A9 both intact and dry cell reaction experiments were carried out. Intact cell reaction was carried out as described previously (Mohamed Ahmed *et al.*, 2009a). The only exception is that at intervals of 0, 30, 60, 90, 120, 150 and 180 min, aliquots of the cell suspension were withdrawn and boiled to stop the reaction. To prepare dried cells, cells suspension was spread on glass Petri dish and dried at room temperature with electric fan. Then the collected cells were further dried in a vacuum desiccator and finely grounded using mortar. The dried cells (75 mg) were suspended in 18 ml of 50 mM potassium phosphate buffer (pH 7.5). The dried cell reaction was started by the addition of homocholine (20 mM) with or without NAD⁺ (5 mM) to the cell suspension. The suspension was incubated on a shaker at 120 rpm and 30°C. At appropriate time intervals (0, 30, 90, 150, 210, and 300 min), aliquots of the cell suspension were withdrawn and boiled for 3–5 min to stop the reaction. After centrifugation, the supernatant was preserved at -20°C and used for metabolites detection as described previously (Mohamed Ahmed *et al.*, 2010). An exception is that TMAPaldehyde concentration in the reaction mixture was quantified by DNPH methods as described elsewhere (Tani *et al.*, 1977). The amount of TMAPaldehyde in the reaction mixture was calculated from the standard curve that generated using different concentration of authentic standard of TMAPaldehyde.

Detection of 3-hydroxypropionate by LC/MS/MS:

Analyses of the intact cell reaction mixtures of *Pseudomonas* sp. strain A9 and the authentic standard of 3-hydroxypropionate were carried out using a Waters LC/MS instrument consisting of a Waters 2695 liquid chromatograph coupled with a Waters Quatromicro API mass spectrometer. LC separations were made using a 4.6 × 150 mm Waters Symmetry C18 column at room temperature and at flow rate of 0.2 ml/min. Solvent A was 0.1% formic acid in acetonitrile, and solvent B was 0.1% formic acid in distilled water. The following gradient of B was applied: 0 min, 100%; 5 min, 100%; 15 min, 0%; 25 min, 0%; 27 min, 100%. MS analysis was done in the negative ESI-MS/MS mode.

RESULTS AND DISCUSSION

Screening for homocholine degrading enzymes activities:

Screening of the homocholine oxidation activities in the isolated strains by replica staining and spectrophotometer assays demonstrated that NAD⁺-dependent dehydrogenase enzymes are predominant in homocholine degrader. In replica staining screening test, about 10 strains showed positive results (formation of purple color), which indicated the presence of NAD⁺-dehydrogenase activities in these isolates (Fig. 1a). Cell free extracts of these isolates again confirmed the presence of both NAD⁺-dependent alcohol and aldehyde dehydrogenase activities (Fig. 1b). Those isolates were chosen as the most superior homocholine degraders and were identified to the specific level according to general principles of microbial classification (Mohamed Ahmed *et al.*, 2009a, b; Mohamed Ahmed *et al.*, 2010). From those isolates, *Pseudomonas* sp. strain A9 was selected as a preferred homocholine degrader, since it showed quite higher activity of both homocholine and TMAPaldehyde dehydrogenases. In the preliminary experiment, these enzymes were found to be unstable during the cell free preparation and against dialysis. Although *Rhodococcus* and *Arthrobacter* strains were also showed both activities but these strains were excluded because cell free extract preparation is quite difficult and need more extraction time that significantly affect the enzyme stability. Moreover, *Rhodococcus* and *Arthrobacter* strains require more cultivation time (48h) compare to *Pseudomonas* strains (24h).

Time course of the growth and enzyme formation of *Pseudomonas* sp. strains A9:

The time course of the formation of homocholine dehydrogenase and cell growth of *Pseudomonas* sp. strain A9 was examined in basal-homocholine medium (Fig. 2). The production of homocholine dehydrogenase activity was significantly increased with the increase in the cell growth (T 660 nm). The maximum enzyme activity formation was observed in late exponential phase at about 24 h. Thereafter, the activity was rapidly decreased after 24h cultivation. Hassan (2008) also found that the formation of TMA-Butanol dehydrogenase activity by *Pseudomonas* sp. 13CM was rapidly decreased after 6 h cultivation. This phenomenon looks similar in quaternary ammonium alcohol degrader. At the moment we do not know the actual reasons for the rapid decrease in the enzyme activity after maximum formation. However, it might be resulted from the inhibition of the cellular enzymes by low molecular weight metabolites that accumulated in high concentration in the culture

medium during the degradation of the substrate and then penetrate into the cellular components. These metabolites might be homocholine analogue such as trimethylamine.




Fig. 1: Screening for dehydrogenases activity in the isolated strains by (A) replica staining and (B) spectrophotometric assays.




Fig. 2: Time course of the growth and homocholine dehydrogenase activity formation of *Pseudomonas* sp. strain A9. Cell-free extract was prepared from cells that grown on 300 ml of culture broth.

Formation of homocholine dehydrogenase on various media:

To assess the induction of homocholine dehydrogenase by *Pseudomonas* sp. strain A9, the bacterium was cultivated on basal media of various homocholine analogues as C and N source, as well as on basal-homocholine medium supplemented with additional C or N sources. The cell free extracts of the above media was prepared and assayed for homocholine dehydrogenase activity. The results (Table 1) showed that homocholine dehydrogenase activity could only be observed in the cell-free extract of cultures grown on homocholine. In assays with cell extracts from cultures grown on choline, 4-*N*-trimethylamino-1-butanol (TMA-Butanol) and glucose, homocholine dehydrogenase activity could not be observed. It was also notable that glucose exerted a total repression of homocholine dehydrogenase activity induction, since no activity was detected on homocholine-glucose growing cells. Similar observation was reported for betaine aldehyde dehydrogenase from *Pseudomonas aeruginosa* (Nagasawa *et al.*, 1976; Velasco-Garcia *et al.*, 2006). Moreover, the intact cell reaction of strain A9 grown on homocholine and glucose was carried out. The formed metabolites TMAPaldehyde, β -alanine betaine and TMA were only detected in the reaction mixture of homocholine growing cells, whereas they were not detected in the reaction mixture of glucose growing cells. These observations demonstrated that the enzymes responsible for degradation of homocholine were induced during the growth on homocholine. The induction of quaternary ammonium compounds degrading activities was also observed by many investigators (Nagasawa *et al.*, 1976; Velasco-Garcia *et al.*, 2006; Setyahadi, 1998).

Table 1: Formation of Homocholine dehydrogenase on various media

Medium (1.0%)	Growth ($\tau_{660\text{nm}}$)	Enzyme activity (mU/ml broth)	Specific activity (mU/mg)
Homocholine	2.57	6.21	30.0
Choline	2.65	0.00	0.0
TMA-Butanol	0.58	0.00	0.0
Glucose plus 0.5% (NH ₄) ₂ SO ₄	3.09	0.00	0.0
Supplemented with carbon source (0.5%)			
Glucose	1.63	0.00	0.0
Glycerol	2.85	3.98	22.2
Supplemented with nitrogen source (0.1%)			
(NH ₄) ₂ SO ₄	3.05	9.34	21.5

Basal medium: 1% substrate, 0.2% K₂HPO₄, 0.2% KH₂PO₄, 0.05% MgSO₄·7H₂O, 0.05% yeast extract, and 0.1% polypeptone.

Stability of homocholine dehydrogenase of *Pseudomonas* sp. strain A9:

The stability of homocholine dehydrogenase was evaluated under several conditions of storage, pH, stabilizers and SH-group protecting reagents in order to optimize the cell free extract preparation and enzyme assay conditions. The overall observation is that the enzyme is unstable and liable to degradation by other enzymes. Addition of protease inhibitor cocktail besides 1mM DTT stabilized the enzyme to some extent, and after dialysis the activity was significantly decreased. Replacement of protease inhibitor cocktail by 1 mM EDTA in the same buffer gave similar results (data not shown), which indicated that metallo-protease might be responsible for the reduction of homocholine dehydrogenase activity. Moreover, the effect of some stabilizers such as DTT (1mM), glutathione (1mM), ethanol (5%) and ethylene glycol (10%) were also tested. The results (Fig. 3) showed that ethanol and ethylene glycol are effective stabilizers of homocholine dehydrogenase activity. About 70% of the enzyme activity was retained after 7 days at 4°C in 50 mM potassium phosphate buffer, pH 7.5, containing either 5% ethanol or 10 % ethylene glycol. Furthermore, different dialysis buffers were examined, and 50 mM potassium phosphate buffer, pH 7.5, gave better results compared to others buffers. On the other hand, the cell free extract was preserved at 4°C without dialysis. The results (Fig. 4) showed that homocholine dehydrogenase retained about 98% of its activity after 3 days. However, after 7 days only 40% of the enzyme activity was retained. This instability of homocholine dehydrogenase makes the purification and characterization of this enzyme is a challenge. In choline degradation pathway, choline dehydrogenase (E.C. 1.1.99.1) is an inner mitochondrial membrane protein that catalyzes the four-electron oxidation of choline to glycine betaine *via* a betaine aldehyde intermediate and requires an electron acceptor other than oxygen. To date, no in depth biochemical or kinetic characterization has been performed on choline dehydrogenase, mainly due to the difficulty in its purification because of the instability of the enzyme *in vitro* (Ghanem, 2006). Recently, Gadda and McAllister- Wilkens (2003) reported the first recombinant choline dehydrogenase. This recombinant form choline dehydrogenase was highly unstable *in vitro*, which hindered any further biochemical and mechanistic investigations of the enzyme (Gadda and McAllister-Wilkens, 2003).




Fig. 3: Effect of stabilizers on the stability of homocholine dehydrogenase of *Pseudomonas* sp. strain A9. Cell free extract (1 ml) was dialyzed overnight against 50 mM potassium phosphate buffer containing either 1mM DTT (■), 1mM glutathione (●), 5% ethanol (▲), or 10% ethylene glycol (◆) and stored at 4°C for 1 week




Fig. 4: Evaluation of the stability of homocholine dehydrogenase during storage at 4°C.

Intact and dry cell reaction of Pseudomonas sp. strain A9 with homocholine:

During the degradation of homocholine by resting cells of *Pseudomonas* sp. strain A9, the amount of homocholine decreased concomitantly with the increase of metabolites, identified as TMAPaldehyde, β -alanine betaine and TMA (Fig. 5). The results confirmed the sequential oxidation of homocholine to TMAPaldehyde and β -alanine betaine. Thereafter, cleavage of C-N bond of β -alanine betaine provided TMA and alkyl chain.

Dry cell reaction was carried out using homocholine (20 mM) as a substrate with and without NAD^+ . The results (Fig. 6a) showed that addition of NAD^+ to the reaction mixture significantly increased the degradation rate of homocholine, as well as the production rate of the metabolites TMA and β -alanine betaine. Whereas, in the reaction mixture without NAD^+ , the degradation rate of homocholine, as well as the metabolites formation was very slow (Fig. 6b). The slight degradation of homocholine in the absence of added NAD^+ might be resulted from the remained NAD^+ with the dried cells. The results demonstrated that the enzymes responsible for the metabolism of homocholine are alcohol and aldehyde dehydrogenases that require NAD^+ as electron acceptor.




Fig. 5: Degradation of homocholine by resting cells of *Pseudomonas* sp. strain A9. Time course degradation of homocholine (■) and the generation of the metabolites, TMAPaldehyde (△) beta-alanine betaine (▲) and TMA (●) by intact cells of *Pseudomonas* sp. strains A9.




Fig. 6: Degradation of homocholine by dried cells of *Pseudomonas* sp. strain A9. Time course degradation of homocholine (■) and the generation of the metabolites beta-alanine betaine (▲) and TMA (●) by dried cells of *Pseudomonas* sp. strains A9. Dry cell reaction was performed with (A) and without (B) NAD⁺.

Substrate specificity of homocholine dehydrogenase:

An attempt was made to determine the substrate spectrum of homocholine dehydrogenase in the crude preparation, because the enzyme is unstable and lost its activity during purification processes. The substrate specificity of homocholine dehydrogenase was determined at 30°C and pH 7.5 using various alcohols and quaternary ammonium compounds in the presence of NAD⁺ as a cofactor. Among alcohols tested, DMA-

Butanol appears to be the most favorable substrate for homocholine dehydrogenase followed by TMA-Butanol, homocholine (TMA-Propanol), 1-butanol, 4-amino-1-butanol and dimethylamino-1-propanol (DMA-Propanol) (Table 2). It is particularly interesting that no detectable activity was found for choline, an analogue of homocholine that is found ubiquitously in nature and the ability to degrade choline are widespread among soil microorganisms. Thus, the ability to degrade homocholine does not go alongside with the ability to catabolize choline. Furthermore, the inability of homocholine dehydrogenase to oxidize choline, in accordance with the induction of the enzyme activity with homocholine, ruled out the enzyme to be a choline-oxidizing enzyme. It was also interesting that homocholine dehydrogenase of strain A9 showed high preference to DMA-Butanol and TMA-Butanol, although this bacterium was unable to grow on these substrates. The inability of the bacterium to grow on these substrates as well as the induction of the enzyme only by homocholine also excluded the enzyme to be a TMA-Butanol-oxidizing enzyme. Recently, a NAD⁺-dependent TMA-Butanol dehydrogenase was purified and characterized from *Pseudomonas* sp. 13CM. This enzyme did not react with homocholine, Dimethyl amino-1- propanol (DMAPropanol), and choline (Hassan, 2008). Thus, it could be assumed that homocholine dehydrogenase is a new enzyme and is not a choline or TMA-Butanol oxidizing enzyme. To ascertain if the activity detected in the cell free extract of homocholine growing cells with different substrate was catalyzed by the same enzyme, crude cell free extract was analyzed by native-PAGE, and then, NAD⁺ dependent activity was located in gels using different substrates. The result showed that only single band of activity was detected (data not shown). The activity band with DMA-Butanol as a substrate was much more intense than that with homocholine as a substrate. This result is in good agreement with the activity measured *in vitro* for both substrates (Table 2). The results obtained demonstrated that homocholine dehydrogenase has a broad substrate range. Broad substrate specificities were also reported for many primary and secondary alcohol and aldehyde dehydrogenases (Vangnai and Arp, 2001; Schenkels and Duine, 2000; Tani *et al.*, 2000; Jaureguibeitia *et al.*, 2007; Jo *et al.*, 2008).

Table 2: Substrate specificity of homocholine dehydrogenase from strain A9


Substrate (33.3 mM)		Activity (mU/ml)	Relative activity (%)
Homocoline	(CH ₃) ₃ N ⁺ (CH ₂) ₂ CH ₂ OH	649	100
TMA-Butanol	(CH ₃) ₃ N ⁺ (CH ₂) ₃ CH ₂ OH	902	139
TMA-Pentanol	(CH ₃) ₃ N ⁺ (CH ₂) ₄ CH ₂ OH	0	0
TMA-Hexanol	(CH ₃) ₃ N ⁺ (CH ₂) ₅ CH ₂ OH	0	0
Choline	(CH ₃) ₃ N ⁺ CH ₂ CH ₂ OH	0	0
β-methyl choline	(CH ₃) ₃ N ⁺ CH(CH ₂)OH	0	0
DMA-Propanol	(CH ₃) ₂ N(CH ₂) ₂ CH ₂ OH	141	22
DMA-Butanol	(CH ₃) ₂ N(CH ₂) ₃ CH ₂ OH	1020	157
DMA-Hexanol	(CH ₃) ₂ N(CH ₂) ₅ CH ₂ OH	0	0
4-Amino-1-butanol	H ₂ N ⁺ (CH ₂) ₃ CH ₂ OH	215	33
1-Butanol	CH ₃ (CH ₂) ₂ CH ₂ OH	236	36
1-Propanol	CH ₃ CH ₂ CH ₂ OH	43	7
2-Propanol	CH ₃ CH (OH) CH ₃	0	0
Ethanol	CH ₃ CH ₂ OH	0	0
Methanol	CH ₃ OH	0	0
L-Carnitine	(CH ₃) ₃ N ⁺ CH ₂ CH(OH)CH ₂ COOH	0	0
D-Carnitine	(CH ₃) ₃ N ⁺ CH ₂ CH(OH)CH ₂ COOH	0	0

Detection of 3-hydroxypropionate dehydrogenase activity:


NAD⁺-dependent 3-hydroxypropionate dehydrogenase activity was detected in the cell free extract of *Pseudomonas* sp. strain A9. This activity was only detected in the cell free extract of homocholine-growing cells and no activity was detected on both choline- and/or glucose-growing cells (Table 3). Furthermore, this enzyme activity was also detected on native-PAGE only in the cell free extract of homocholine growing cells (Fig. 7a). These results demonstrated the induction of 3-hydroxypropionate dehydrogenase activity by homocholine and indicated the presence of 3-hydroxypropionate as an intermediate metabolite in the degradation pathway of homocholine by *Pseudomonas* sp. strain A9. Moreover, to confirm the presence of 3-hydroxypropionate as an intermediate metabolite, the intact cell reaction mixtures of strain A9 were analyzed by LC/LC-MS. The results demonstrated the accumulation of 3-hydroxypropionate as an intermediate metabolite (Fig. 7b). The mass spectrum (M⁺ 90.08) and the retention time (6.7 min) of the observed metabolite, agreed with those of authentic standard of 3-hydroxypropionate. In a similar study, cleavage of C-N bond of choline by *Candida tropicalis* was accompanied by formation of trimethylamine and ethylene glycol (Mori *et al.*, 1988). 3-Hydroxypropionate is of special interest in view of that the biodegradable polymers of it can potentially replace lot kinds of traditional petrochemistry-based polymers and be used in some new fields such as surgical biocomposite materials and drug release material (Jiang *et al.*, 2009).

Table 3: NAD⁺-dependent 3-hydroxypropionate dehydrogenase activity in the cell extracts of homocholine-, choline- and glucose-growing cells

Medium Substrate (1.0%)	Activity (mU/ml)
Homocholine	125
Choline	0
Glucose	0

**Fig. 7:** (A) Non-denaturing gel (7.5% gel) stained for NAD⁺-dependent 3-hydroxypropionate dehydrogenase activity. Lane 1; homocholine-growing cells, lane 2; choline-growing cells, and lane 3; glucose-growing cells. (B) LC/MS/MS analysis of 3-hydroxypropionate in the degradation pathway of homocholine by the intact cells of *Pseudomonas* sp. Strain A9. Intact cell reaction mixture of strain A9 (upper) and authentic standard of 3-hydroxypropionate (lower).**Conclusion:**

Overall, it could be concluded that homocholine degrading enzymes are widespread in aerobic microorganisms such as *Arthrobacter*, *Rhodococcus* and *Pseudomonas*. In *Pseudomonas* sp. strain A9, homocholine was oxidized to TMAPaldehyde by homocholine dehydrogenase, and consequently, TMAPaldehyde oxidized to β -alanine betaine by a NAD⁺-dependent aldehyde dehydrogenase. Thereafter, cleavage of β -alanine betaine C-N bond yielded trimethylamine and 3-hydroxypropionate (C-3 moiety). 3-Hydroxypropionate was further oxidized to malonate semi-aldehyde by a NAD⁺-dependent 3-hydroxypropionate dehydrogenase (Fig. 8). Further studies shall specifically focus on the purification and characterization of homocholine degrading enzymes.

**Fig. 8:** Proposed degradation pathway of homocholine by *Pseudomonas* sp. strain A9

REFERENCES

- Boksa, P., and B. Collier, 1980. Acetylation of homocholine by rat brain: subcellular distribution of acetylhomocholine and studies on the ability of homocholine to serve as substrate for choline acetyltransferase in situ and in vitro. *J Neurochem*, 34(6): 1470-1482.
- Carroll, P.T., and J.M. Aspry, 1980. Subcellular origin of cholinergic transmitter release from mouse brain. *Science*, 210: 641-642.
- Cavener, D.R., 1992. GMC oxidoreductases. A newly defined family of homologous proteins with diverse catalytic activities. *J Mol Biol.*, 223: 811-814.
- Channon, H.J., A.P. Platt, and J.A.B. Smith, 1937. The dietary prevention of fatty livers. Two analogues of choline. *Biochem J*, 31: 1736-1742.
- Collier, B., S. Lovat, D. Ilson, L.A. Barker and T.W. Mittag, 1977. The uptake, metabolisms and release of homocholine: studies with rat brain synaptosomes and cat superior cervical ganglion. *J Neurochem*, 28: 331-339.
- Gadda, G., and E.E. McAllister-Wilkins, 2003. Cloning, Expression and Purification of Choline Dehydrogenase from the Moderate Halophile *Halomonas elongata*. *Appl Environ Microb*, 69: 2126-32.
- Ghanem, M., 2006. On the Mechanistic Roles of the Protein Positive Charge Close to the N(1) Flavin Locus in Choline Oxidase. Ph. D. Thesis, Georgia State University.
- Hassan, M., 2008. Enzymatic studies on the degradation of 4-N-trimethylamino-1-butanol by *Pseudomonas* sp. 13CM. Ph. D. Thesis, Tottori University, Japan.
- Jaureguibeitia, A., L. Saa, M.J. Llama, and J.L. Serra, 2007. Purification, characterization and cloning of aldehyde dehydrogenase from *Rhodococcus erythropolis* UPV-1. *Appl Microbiol Biotechnol*, 73: 1073-1086.
- Jiang, X., X. Meng, and M. Xian, 2009. Biosynthetic pathway for 3-hydroxypropionic acid production. *Appl Microbiol Biotechnol.*, 82: 995-1003.
- Jo, J.-E., S.M. Raj, C. Rathnasingh, E. Selvakumar, W.-C. Jung and S. Park, 2008. Cloning, expression, and characterization of an aldehyde dehydrogenase from *Escherichia coli* K-12 that utilizes 3-hydroxypropionaldehyde as substrate. *Appl Microbiol Biotechnol.*, 81: 51-60.
- Johannes, T.W., R.D. Woodyer and H. Zhao, 2006. High-throughput screening methods developed for oxidoreductase. In: Reymond J-L, *Enzyme Assays: High-throughput Screening, Genetic Selection and Fingerprinting*. WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, pp: 77-93.
- Mohamed Ahmed, I.A., J. Arima, T. Ichianagi, E. Sakuno and N. Mori, 2009a. Isolation and characterization of 3-N-trimethylamino-1-propanol degrading *Rhodococcus* sp. strain A2. *FEMS Microbiol lett*, 296(2): 219-225.
- Mohamed Ahmed, I.A., J. Arima, T. Ichianagi, E. Sakuno, and N. Mori, 2009b. Isolation and characterization of 3-N-trimethylamino-1-propanol degrading *Arthrobacter* sp. strain E5. *Res J Microbiol.*, 4(2): 49-58.
- Mohamed Ahmed, I.A., J. Arima, T. Ichianagi, E. Sakuno and N. Mori, 2010. Isolation and characterization of homocholine degrading *Pseudomonas* sp. strain A9 and B9b. *World J Microbiol Biotechnol.*, 26: 1455-1464.
- Mori, N., S. Fuchigami, and Y. Kitamoto, 2002. Purification and properties of betaine aldehyde dehydrogenase with high affinity for NADP from *Arthrobacter globiformis*. *J Biosci Bioeng*, 93: 130-135.
- Mori, N., B. Kawakami, K. Hyakutome, Y. Tani and Y. Yamada, 1980. Characterization of betaine aldehyde dehydrogenase from *Cylindrocarpus didymum* M-1. *Agric Biol Chem*, 44: 3015-3016.
- Mori, N., K. Shirakawa, K. Uzura, Y. Kitamoto, and Y. Ichikawa, 1988. Formation of ethylene glycol and trimethylamine from choline by *Candida tropicalis*. *FEMS Microbiol Lett*, 51: 41-44.
- Mori, N., N. Yoshida, and Y. Kitamoto, 1992. Purification and properties of betaine aldehyde dehydrogenase from *Xanthomonas translucens*. *J Ferment Bioeng*, 73: 352-356.
- Nagasawa, T., N. Mori, Y. Tani and K. Ogata, 1976. Characterization of choline dehydrogenase of *Pseudomonas aeruginosa* A-16. *Agric Biolog Chem*, 40: 2077-2084.
- Russell, B.L., B. Rathinasabapathi and A.D. Hanson, 1998. Osmotic stress induces expression of choline monooxygenase in sugar beet and amaranth. *Plant Physiol.*, 116: 859-865.
- Schenkels, P., and J. A. Duine, 2000. Nicotinoprotein (NADH-containing) alcohol dehydrogenase from *Rhodococcus erythropolis* DSM 1069: an efficient catalyst for coenzyme-independent oxidation of a broad spectrum of alcohols and the interconversion of alcohols and aldehydes. *Microbiol.*, 146: 775-785.
- Schmid, A., J.S. Dordick, B. Hauer, A. Kiener, M. Wubbolts, and B. Witholt, 2001. Industrial biocatalysis today and tomorrow. *Nature*, 409: 258-268.
- Setyahadi, S., 1998. Structure and Function of D-carnitine dehydrogenase of *Agrobacterium* sp. 525a and enzymatic production of L-carnitine. Ph.D thesis, Tottori University, Tottori, Japan.

Tani, A., Y. Sakai, T. Ishige, and N. Kato, 2000. Thermostable NADP⁺-dependent medium-chain alcohol dehydrogenase from *Acinetobacter* sp. strain M-1: Purification and characterization and gene expression in *Escherichia coli*. *Appl Environ Microbiol.*, 66(12): 5231-5235.

Tani, Y., N. Mori, and K. Ogata, 1977. A new fungal enzyme, choline oxidase. *Agric Biol Chem*, 41: 1101-1102.

Vangnai, A.S., and D.J. Arp, 2001. An inducible 1-butanol dehydrogenase, a quinohaemoprotein, is involved in the oxidation of butane by *Pseudomonas butanovora*. *Microbiol.*, 147: 745-756.

Velasco-Garcia, R., M.A. Villalobos, M.A. Ramirez-Romero, C. Mujica-Jimenez, G. Iturriaga and R.A. Munoz-Clares, 2006. Betaine aldehyde dehydrogenase from *Pseudomonas aeruginosa*: cloning, over-expression in *E. coli* and regulation by choline and salt. *Arch Microbiol.*, 185: 14-22.